

# Isokinetic muscle strength and readiness to return to sport following anterior cruciate ligament reconstruction: is there an association? A systematic review and a protocol recommendation

Marit Baste Undheim,<sup>1</sup> Ciaran Cosgrave,<sup>1</sup> Enda King,<sup>1,2</sup> Siobhán Strike,<sup>2</sup> Brendan Marshall,<sup>1,3</sup> Éanna Falvey,<sup>1,4</sup> Andrew Franklyn-Miller<sup>1,5</sup>

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/bjsports-2014-093962>).

<sup>1</sup>Sports Medicine Department, Sports Surgery Clinic, Dublin, Ireland

<sup>2</sup>Department of Life Sciences, Roehampton University, London, UK

<sup>3</sup>Insight Centre for Data Analytics, Dublin City University, Dublin, Ireland

<sup>4</sup>Department of Medicine, University College Cork, Cork, Ireland

<sup>5</sup>Centre for Health, Exercise and Sports Medicine, University of Melbourne, Melbourne, Australia

## Correspondence to

Marit Baste Undheim, Sports Medicine Department, Sports Surgery Clinic, Santry Demesne, Dublin 9, Ireland; [mundheim@gmail.com](mailto:mundheim@gmail.com)

Accepted 1 June 2015

## ABSTRACT

**Introduction** Following anterior cruciate ligament reconstruction (ACLR), strength is a key variable in regaining full function of the knee. Isokinetic strength is commonly used as part of the return to sport (RTS) criteria.

**Aim** We systematically reviewed the isokinetic strength evaluation protocols that are currently being used following ACLR. A secondary aim was to suggest an isokinetic protocol that could meet RTS criteria.

**Method** Articles were searched using ScienceDirect, PubMed and Sage Journals Online, combined with cross-checked reference lists of the publications. Protocol data and outcome measurements and RTS criteria were extracted from each article included in the review.

**Results** 39 studies met the inclusion criteria and reported their isokinetic strength evaluation protocol following ACLR. The variables that were most commonly used were concentric/concentric mode of contraction (31 studies), angular velocity of 60°/s (29 studies), 3–5 repetitions (24 studies), range of motion of 0–90° (6 studies), and using gravity correction (9 studies). 8 studies reported strength limb symmetry index scores as part of their RTS criteria.

**Conclusions** There was no standardised isokinetic protocol following ACLR; isokinetic strength measures have not been validated as useful predictors of successful RTS. We propose a standard protocol to allow consistency of testing and accurate comparison of future research.

## INTRODUCTION

Injury to the anterior cruciate ligament (ACL) is a common acute knee injury in sports and it provides many challenges to clinicians and researchers.<sup>1</sup> ACL reconstruction (ACLR) surgery is common among athletes who want to return to sport (RTS).<sup>2–4</sup> After ACLR, athletes have demonstrated muscle strength deficit, decreased stability and force attenuation for up to 2 years after reconstruction<sup>3 5–9</sup> and this may influence future knee injury risk.<sup>10 11</sup>

In total 6 to 27% of athletes who RTS after ACLR sustain a new ACL injury (to the same or opposite knee) within 10 years postsurgery.<sup>12–15</sup> As muscle strength deficit has been associated with the potential risk of future knee injury, increasing muscle strength of the quadriceps and hamstrings is a key factor for successful RTS after ACLR surgery.<sup>16–19</sup> Strength evaluation tests following ACLR are commonly performed as part of a

battery of tests in clinical and research settings.<sup>20 21</sup> Isokinetic dynamometry provides an objective measure of muscle strength and is used in sport, research and clinical settings.<sup>20 21</sup> A criticism of isokinetic dynamometry is that it lacks functional relevance to sporting and training situations. However, it is considered the ‘gold standard’ for measuring muscle strength<sup>22</sup> and its convenience, reproducibility and reliability support its use as an appropriate method of assessment after ACLR.<sup>22–24</sup> Two key aspects to indicate knee strength following ACLR surgery are limb symmetry index of the injured and uninjured leg, and hamstring to quadriceps (H/Q) ratios of the injured leg. These measures have commonly been referred to and used as part of the RTS criteria.<sup>2 17 25</sup>

Part of successful rehabilitation is to return the patient to ‘normal’ strength. ‘Normal’ limb symmetry index values are reported to be between >70–90%,<sup>2 26 27</sup> while a ‘normal’ H/Q ratio has been reported to be between 0.5 and 0.8.<sup>28 29</sup> Many clinicians, therefore, aim to return the patient back to these values and use these values as part of their RTS criteria.<sup>28</sup> However, the torque production and results may be affected by aspects of the isokinetic protocol<sup>16</sup> (including modes of contraction, angular velocity, range of motion, number of repetitions and gravity correction).<sup>30–32</sup>

The protocols used for isokinetic strength evaluation following ACLR are, therefore, important factors to review when considering an athlete’s readiness to RTS. The primary aim of this study was to systematically review and report the isokinetic strength evaluation protocols that are currently in use when assessing the muscle strength of patients who have undergone ACLR. The secondary aim is to suggest an appropriate strength evaluation protocol following ACLR in accordance with the RTS criteria.

## METHOD

### Search strategy

Two authors (MU, CC) conducted a systematic literature search of the ScienceDirect, PubMed, and Sage Journals Online databases, combined with cross-checked reference lists of the publications using PRISMA guidelines.<sup>33</sup> The literature search was performed between October and December 2013. Two main keywords, “anterior cruciate ligament reconstruction” and “isokinetic dynamometer”, were used to search for articles. In each

**To cite:** Undheim MB, Cosgrave C, King E, et al. *Br J Sports Med* Published Online First: [please include Day Month Year] doi:10.1136/bjsports-2014-093962

## Review

search, the two main keywords were followed by either “muscle strength” or “peak torque” or “limb asymmetry” or “return to sport”.

### Selection criteria

The selection criteria were determined by the lead author (MU) before commencing the article search. To be included in this review, the articles had to have (A) participants who had undergone ACLR surgery; (B) evaluated knee flexor and extensor isokinetic muscle strength up to 24 months following ACLR; (C) been published between January 1980 and October 2013 and (D) been published in English. Studies that assessed participants who had undergone ACLR revision or multiple-ligament reconstruction were excluded. If an article evaluated knee flexors or knee extensors only, these were excluded. Testing of both knee extensors and flexors were included, as both quadriceps and hamstrings have been reported to play a crucial role in the knee function following ACLR<sup>5 6 8</sup> and both are reported as the most commonly used graft-harvesting sites for ACLR.<sup>3 5–9</sup> Articles that evaluated strength at more than 24 months following ACLR were excluded, as strength deficiencies are seen up to 24 months following ACLR.<sup>3 5 6 9</sup> Articles were also excluded if isokinetic muscle strength was tested lying down rather than when seated, and if data were repeated in another article already included in the review.

The title and abstract of each study were reviewed first and if it was not clear whether a study was appropriate for inclusion, the full text of the article was examined. The selection criteria were applied by two independent reviewers (MU, CC). Consensus was used to resolve any disagreements between reviewers, with a third reviewer (BM) consulted if consensus was not achieved.

### Data extraction and analysis

Data was extracted from included papers by the lead author (MU). Data identified for descriptive analysis included the isokinetic dynamometer protocol and isokinetic outcome measures. Specifically, data pertained to modes of contraction, angular velocity, number of repetitions, set range of motion and if gravity correction was used. The outcome measurements were peak torque and total work, and the limb symmetry index, H/Q ratio and/or magnitude of the strength outcome measure. Articles that reported RTS criteria in their rehabilitation protocol or reported recommended values for RTS in their article were included as the second aim of the review. The limb symmetry index values and H/Q ratios were extracted for these articles. These values were then reviewed and related to the strength evaluation protocols.

### Assessment for risk of bias

The studies included in the systematic review were of different methodological design. The Downs and Black<sup>34</sup> checklist was used for assessing methodological quality because it is appropriate for determining the quality of both randomised and non-randomised studies.<sup>35</sup> The Downs and Black<sup>34</sup> checklist comprises of 27 items and a maximum score of 32 can be obtained if all criteria are met. The checklist is divided into five sections where each article is assessed on its reporting, external validity, internal validity -bias, internal validity—confounding (selection bias) and power. The articles were divided into quality-level categories according to previous literature:<sup>36</sup> excellent (26–28), good (20–25), fair (15–19) and poor ( $\leq 14$ ). The assessment of methodological quality was completed by the same two reviewers (MU, CC), independently.

## RESULTS


A total of 548 published studies were identified in the original search of the databases and other resources. After duplicates were removed, 263 studies were screened through titles and abstracts. Seventy-five remaining studies were reviewed in full text and 36 of these studies were excluded: 15 studies evaluated strength at more than 24 months following ACLR, 8 studies evaluated knee flexors only, 9 evaluated knee extensors only, 1 study repeated data used in a previous study, 1 study evaluated strength lying down and 2 studies evaluated isometric strength only. A final yield of 39 studies<sup>3 6–9 21 23 29 37–67</sup> were included in this systematic review as presented in the flow chart (figure 1).

Methodological quality assessment scores ranged between scores of 12 and 26 of a possible 32 points (see online supplementary table S1), with a median score of 18. None of the papers were classified as excellent. Seven were classified as good,<sup>43 45–47 53 55 66 26</sup> as fair<sup>3 5–9 21 23 37–39 41 42 44 48–50 52 54 56 60–67</sup> and 6 as poor.<sup>29 40 51 57–59</sup> Thirty studies did not include all important adverse events that could have an effect on the intervention reported (criterion 8),<sup>5–7 9 21 23 29 38–40 42–52 54 56 57 59 60 63 64–67</sup> 23 studies did not adequately describe the source population and how they were recruited (criterion 11 and 12),<sup>3 5 6 8 21 37–40 42 44 48–52 54 57–60 64 65</sup> 24 studies did not describe if the source population was recruited in the same time period (criterion 22),<sup>3 8 9 23 29 37–42 44 48–52 54–58 63 64 66</sup> and 29 studies did not report sample size calculation (criterion 27).<sup>3 6–8 21 23 29 38–44 49–53 56–61 63–67</sup> Other criterion, such as criterion 22 and 23, were not commonly fulfilled due to most studies not being randomised, while criterion 14 and 15 were not commonly reported due to lack of blinding.

Variables extracted for each study can be found in online supplementary table S2. Table 1 shows the number and percentage of studies reporting the protocol variables. All studies reported mode of contraction as concentric/concentric, where a patient completes concentric knee extension and flexion contractions, or concentric/eccentric, where a patient performs concentric knee extension and eccentric knee flexion. Mode of contraction and angular velocity were reported in all the studies reviewed, but not all studies reported all aspects of their protocols. Of the 39 studies, 11 (28%) did not report number of repetitions,<sup>7 9 39 43 45 46 52 54 58 61 66</sup> 25 (64%) did not report range of motion,<sup>3 6–9 29 38 39 41 43 47–51 54 55 57 58 60 61 63 64–66</sup> and 29 (74%) studies did not report if gravity correction was used.<sup>3 6 7 9 21 29 37 38–45 47 49 50 53–55 57 58 60 61 64–67</sup>

In table 2, the outcome measurements and results that were reported are presented. Peak torque was the most commonly reported, with 32 of 38 studies reporting it as one of the outcome measurements. One study did not report the outcome measure that they used.<sup>41</sup> All 39 studies reported their results as a limb symmetry index score, a H/Q ratio and/or a magnitude. Five (13%) reported magnitude only, that is, the studies only reported strength measures in their respective outcome measurement (torque and/or work) and did not compare injured and uninjured leg strength measures or H/Q ratios. Thirty-four studies (87%) reported their results as limb symmetry index and/or H/Q ratio. All 34 studies reported limb symmetry index scores, while only 6 (18%) reported H/Q ratio.

Fifteen studies (38%) reported strength as a component of their RTS criteria. Of these, eight studies used limb symmetry index as the RTS criterion. The limb symmetry index scores are reported in table 2, and show a variation from >70–90%. The remaining seven of these studies reported strength criteria as



**Figure 1** Preferred reporting items for systematic reviews and meta-analysis (PRISMA) flow chart of search and included and excluded studies.

returning to ‘normal’/‘adequate’/‘good’ or preoperation level of strength, rather than a specific score.

From 39 studies, 8 (21%) reported objective RTS strength criteria.<sup>40 41 43 50 51 54 56 66</sup> The protocol data, outcome measurements and RTS criteria for these studies are presented in [table 3](#). All RTS criteria were reported as limb symmetry index scores.

## DISCUSSION

We found no standardised isokinetic strength evaluation protocol following ACLR. If there is a need to objectively assess a patient’s strength following ACLR, a requirement exists for a defined protocol. In this systematic review, mode of contraction, angular velocity, number of repetitions, range of motion and the use of gravity correction were the main variables that were extracted from each article.

### Isokinetic strength evaluation protocol

Thirty-one studies (79%) used concentric mode of contraction for testing rather than using concentric and eccentric contractions. The mode of contraction significantly affects torque output. Eccentric torque output is significantly greater than concentric torque output and could, therefore, be misleading if

indirectly compared.<sup>68</sup> In sports, concentric and eccentric contractions are performed, and therefore post-ACLR, testing concentric and eccentric strength could be important.

Previous research has reported that patients find the eccentric movement more difficult than the concentric movement<sup>69 70</sup> as it requires greater skill and motor control compared with concentric contraction, and as such may yield lower reliability and reproducibility.<sup>69</sup> Previous studies have shown a ‘very high’ and ‘high’ reproducibility and reliability for isokinetic concentric and eccentric knee extension and flexion; however, in all studies, the eccentric knee extension and flexion reliability and reproducibility is lower than the concentric.<sup>24 69 70 71</sup> Although the concentric mode of contraction is more commonly used in clinical and research settings,<sup>40 59 66</sup> it does not necessarily mean it is the better mode of contraction. However, due to its frequency of use, there is an abundance of comparable data.<sup>3 6 7 9 21 29 37–41 43 44 48–51 53–62 64–67</sup> If eccentric strength evaluation is used, a familiarisation session may be required to ensure good reliability and different RTS criteria may have to be considered.

Angular velocity influences both the torque and work output, and as such, should be considered when assessing strength and using it as part of the RTS criteria.<sup>30 40 56 59</sup> The two most common angular velocities reported were 60°/s and 180°/s in

## Review

**Table 1** Strength evaluation protocol data used in studies (N (%))

| Protocol                | Studies (N (%)) |
|-------------------------|-----------------|
| Mode of contraction     |                 |
| Concentric/concentric   | 31 (79)         |
| Concentric/eccentric    | 8 (21)          |
| Angular velocity (°/s)* |                 |
| 30                      | 1 (3)           |
| 60                      | 29 (74)         |
| 90                      | 3 (8)           |
| 120                     | 5 (13)          |
| 180                     | 18 (46)         |
| 230                     | 1 (3)           |
| 240                     | 11 (28)         |
| 270                     | 1 (3)           |
| 300                     | 7 (18)          |
| 450                     | 1 (3)           |
| Number of repetitions*  |                 |
| 3–5                     | 24 (62)         |
| 6–10                    | 7 (18)          |
| >10                     | 7 (18)          |
| Not reported            | 11 (28)         |
| Range of motion (°)*    |                 |
| 0–100                   | 3 (8)           |
| 0–90                    | 6 (15)          |
| 5–90                    | 1 (3)           |
| 10–90                   | 3 (8)           |
| 20–90                   | 2 (5)           |
| 40–90                   | 2 (5)           |
| 45–90                   | 1 (3)           |
| Not reported            | 25 (64)         |
| Gravity correction      |                 |
| Yes                     | 9 (23)          |
| No                      | 1 (3)           |
| Not reported            | 29 (74)         |

\*Some studies used more than one test, therefore the total sum of percentages do not equal 100%.

29 (74%) and 18 (46%) studies, respectively. Abundant data exists for these selected velocities and have been shown to highlight strength deficits.<sup>37 40 50 55 57 61 63 66</sup> Torque output decreases as angular velocity increases above 60°/s, and the maximum torque output is shown to be between 0 and 60°/s.<sup>72</sup> For this reason, if the angular velocity that is used for strength evaluation is greater than 60°/s, strength deficiencies might not be highlighted. If strength deficiencies are not highlighted due to high angular velocity, it might be falsely assumed that the RTS strength criterion is achieved. It is, therefore, important to use an angular velocity that will consistently highlight strength deficiencies.

Other strength evaluation protocol data, such as number of repetitions, range of motion and the use of gravity correction, will also affect the outcome measures and depend largely on the previous protocol variables used. When reporting peak torque, five repetitions are recommended because peak torque is achieved at the fourth repetition.<sup>73</sup> By using five repetitions, there is a greater chance of recording the highest peak torque and less chance of fatigue. Maximal range of motion is recommended to allow the patient to maximise the isokinetic phase by reaching the preset angular velocity and achieve maximum voluntary activation.<sup>74</sup> Angular position can provide valuable information about the mechanical properties of the contracting

**Table 2** Outcome measurements and results reported in studies (N (%))

| Result measures                   | Studies (N (%)) |
|-----------------------------------|-----------------|
| Outcome measures*                 |                 |
| Peak torque                       | 32 (82)         |
| Total work                        | 11 (28)         |
| Peak torque and total work        | 9 (23)          |
| Not reported                      | 1 (3)           |
| Results reported as*              |                 |
| LSI and/or H/Q ratio              | 34 (87)         |
| Magnitude of outcome measure only | 5 (13)          |
| RTS strength criterion*           |                 |
| LSI: (%)                          |                 |
| ≥70–75                            | 4 (10)          |
| ≥80–85                            | 3 (8)           |
| ≥90–95                            | 2 (5)           |
| Not reported                      | 31 (79)         |

\*Some studies report more than one variable for the results, therefore the total sum of percentages do not equal 100%.

H/Q, Hamstring to Quadriceps; LSI, Limb Symmetry Index; RTS, return to sport.

muscle and range of motion can affect the peak torque output; therefore, it is important to set the same range of motion for each patient.<sup>72</sup> Finally, the use of gravity correction is important for reporting accurate strength results for the H/Q muscles.<sup>11</sup> If gravity correction is not used during concentric/concentric mode of testing, the quadriceps strength scores will decrease due to the upward motion acting against gravity and hamstrings strength scores will increase due to the downward motion acting with gravity.<sup>75</sup>

### Reporting outcome measures and results

Conventionally, isokinetic strength measures are reported as peak torque and/or total work.<sup>20</sup> Peak torque during isokinetic movements is a measure of the maximal force exerted during knee extension and flexion. Since it eliminates any submaximal repetitions, peak torque is a good measure of maximal strength.<sup>76</sup> Conversely, total work is a better measure of the endurance of the muscle.<sup>52 72</sup> In this review, peak torque was reported in 32 studies and total work was reported in 11 studies. The limb symmetry index scores in the studies reporting-specific RTS strength criteria were all reported using peak torque. Both peak torque and total work have shown to have high test-retest reliability and to be accurate measures of strength for knee extensors and flexors.<sup>20 69 71 77</sup> As peak torque is the most commonly used outcome measure when reporting limb symmetry index and H/Q ratio, an abundance of comparable data is available.<sup>6 9 43 46 47 51–54 59</sup> Peak torque should, therefore, be used as an outcome measure when considering the RTS strength criteria, though the importance of quadriceps and hamstring muscle endurance following ACLR and its role as part of RTS criteria should be further investigated. The limb symmetry index and H/Q ratio are dependent on the method of testing, and a change in the protocol variables, such as mode of contraction or angular velocity, will change the outcome measures and in turn, the results reported.<sup>30–32 73</sup>

### RTS strength criteria

A secondary finding of the research was that although isokinetic strength is frequently measured in ACLR research, only 15 studies (38%) reported isokinetic strength as part of their RTS



**Table 3** Isokinetic muscle strength evaluation protocol data, outcome measurements and results used in studies reporting RTS strength criteria

| Study                                | Time of strength assessment postsurgery (months) | Mode of contraction | Angular velocity (°/s) | Gravity correction | Repetitions  | Range of motion (°)                   | Outcome measurements    | Isokinetic results reported as: | Isokinetic strength LSI RTS criteria (%)               |
|--------------------------------------|--|---------------------|------------------------|--------------------|--------------|---------------------------------------|-------------------------|---------------------------------|--|
| Cardone <i>et al</i> <sup>40</sup>   | 2, 4, 6  | Conc/conc           | 60, 180, 240           | Not reported       | 4            | 20–90 (at 2 months postsurgery), 0–90 | Peak torque, total work | LSI, magnitude                  | ≥75  |
| Carter and Edinger <sup>41</sup>     | 6  | Conc/conc           | 180, 300               | Not reported       | 5, 15        | Not reported                          | Not reported            | LSI                             | ≥80  |
| Feller and Webster <sup>43</sup>     | 4, 8, 12   | Conc/conc           | 60, 240                | Not reported       | Not reported | Not reported                          | Peak torque             | LSI                             | ≥70  |
| Keays <i>et al</i> <sup>50</sup>     | 6  | Conc/conc           | 60, 120                | Not reported       | 5            | Not reported                          | Peak torque             | LSI, magnitude                  | ≥90  |
| Kobayashi <i>et al</i> <sup>51</sup> | 1, 6, 12, 24                                     | Conc/conc           | 60, 180                | Yes                | 5            | Not reported                          | Peak torque             | LSI                             | ≥70 (quadriceps)<br>≥85 (quadriceps), ≥90% (hamstring) |
| Osteras <i>et al</i> <sup>54</sup>   | 6  | Conc/conc           | 60, 240                | Not reported       | Not reported | Not reported                          | Peak torque             | LSI, magnitude                  | ≥85 (quadriceps)                                       |
| Silva <i>et al</i> <sup>56</sup>     | 5  | Conc/conc           | 60, 180                | Yes                | 3            | 0–90                                  | Peak torque             | LSI, H/Q ratio, magnitude       | ≥85 (quadriceps)                                       |
| Yuksel <i>et al</i> <sup>66</sup>    | 18–24  | Conc/conc           | 60, 180                | Not reported       | Not reported | Not reported                          | Peak torque             | LSI                             | >70  |

Conc, Concentric; H/Q, Hamstring to Quadriceps; LSI, Limb symmetry index; RTS, return to sport.

criteria. Of these 15 studies, 8 reported RTS isokinetic strength criteria, such as limb symmetry of the quadriceps and hamstring strength, but different desirable ranges of limb symmetry index were reported in these studies, indicating the variability in the RTS strength criteria being used.<sup>40 41 43 50 51 54 56 66</sup> Previous research has reported that the limb symmetry index varies between concentric and eccentric contractions.<sup>47 52 63</sup> These studies do not show if one mode of contraction highlights side-to-side differences more than the other and it might be beneficial to use both modes of contractions as it may highlight any asymmetries between the injured and uninjured leg. Seven of the studies used 60°/s as their lowest angular velocity, while one used 180°/s. The studies reported a desirable limb symmetry index that ranged from ≥70% to 90%. However, many studies report an increase in limb symmetry index with higher angular velocities.<sup>7 29 30 45 59</sup> This could mean that in some studies the RTS criterion was met more easily than in others.

'Normal' strength has been classified in relation to the strength of the uninjured limb; thus limb symmetry index scores are commonly reported following ACLR.<sup>41</sup> In this review, 34 (87%) studies reported their results as limb symmetry index scores and it was the only result included as a measure for strength RTS criterion. Barber-Westin and Noyes<sup>16</sup> concluded that a limb symmetry index of >90% should be reached before RTS, while Van Grinsven *et al*<sup>19</sup> reported that a limb symmetry index of >95% was required. In spite of this, of the eight studies<sup>40 41 43 50 51 54 56 66</sup> that reported limb symmetry values as part of their RTS criteria, only Cardone *et al*<sup>40</sup> reported if their RTS strength criterion predicted RTS. Their participants did not show any signs of difficulty on RTS, and they, therefore, suggested that a deficit of 25% in extension strength between injured and non-injured leg at an angular velocity of 60°/s is a reliable RTS criterion. This study, however, had a relatively short follow-up and did not extend its conclusion to hamstring strength. It cannot be assumed that any of these RTS strength criteria are reliable. Together, these findings highlight that isokinetic knee strength has not been sufficiently validated as a useful criterion measure for RTS.

### Methodological considerations

A limitation of the current review is that due to the historical 33 year span of data (1980–2013), changes in surgical and rehabilitation practices (arthroscopic surgery, immediate weight-bearing, accelerated rehabilitation protocols) have occurred. These likely affected the isokinetic dynamometer protocols such as the specific angular velocity and range of motion protocols applied. Furthermore, inclusion criteria did not specify a minimum period post-ACLR at which the strength assessment occurred. However, studies which had a short time line post-ACLR testing, also tested at a time period of at least 5 months postsurgery.<sup>7 29 40 43 45–47 49 51 56 57</sup> Previous research has demonstrated intermanufacturer (Cybex, Kin-Com and Biodex) variance exists using the same protocol.<sup>78 79</sup> This highlights a further potential limitation that the review did not take into account the different types of isokinetic dynamometers utilised. Following the methodological quality assessment, a potential limitation was the studies' poor description of the source population and recruitment. One could hypothesise that this could affect the generalisability of the results. A final limitation is that the prospective relationship between strength measures and RTS outcomes has yet to be determined. Therefore, the secondary aim of suggesting an isokinetic strength protocol based on the RTS criteria was not possible to fulfil. Instead, we

## Review

propose an isokinetic strength protocol based on those proposed in the current literature.

In summary, we propose the following protocol as the most valid assessment for RTS: five repetitions of concentric knee extension and flexion at an angular velocity of 60°/sec, at a set range of motion of 0–100°, using gravity correction, and measuring peak torque.

## What are the new findings?

- ▶ The literature does not show a clear standardised strength evaluation protocol following anterior cruciate ligament reconstruction (ACLR).
- ▶ There is no consensus in the literature as to what is an appropriate return to sport (RTS) strength criteria following ACLR.
- ▶ The proposed standardised isokinetic strength protocol following ACLR includes five repetitions of concentric knee extension and flexion at 60°/s.

## How might it impact on clinical practice in the near future?

- ▶ The proposed standard protocol can be used in prospective studies to analyse if isokinetic strength is associated with successful return to sport (RTS).
- ▶ It will standardise isokinetic assessment of the knee post-ACLR in clinical practice and enable effective comparison of data and the potential to share this across clinics.
- ▶ It may contribute to RTS criteria following ACLR.

**Twitter** Follow Marit Undheim at @maritundheim and Enda King at @enda\_king

**Contributors** MU, CC, EK and AF-M contributed to the conception and design of this review. MU and CC reviewed the studies included in this systematic review. MU, EK, BM and SS drafted the article, while all authors revised it critically for important intellectual content. All authors approved the final version of the manuscript. MU is responsible for the overall content of the manuscript.

**Competing interests** None declared.

**Provenance and peer review** Not commissioned; externally peer reviewed.

## REFERENCES

- 1 Renström PA. Eight clinical conundrums relating to anterior cruciate ligament (ACL) injury in sport: recent evidence and a personal reflection. *Br J Sports Med* 2023;47:367–72.
- 2 Beynon BD, Johnson RJ, Abate JA, et al. Treatment of anterior cruciate ligament injuries, Part I. *Am J Sports Med* 2005;33:1579–602.
- 3 Aune AK, Holm I, Risberg MA, et al. Four-strand hamstring tendon autograft compared with patellar tendon-bone autograft for anterior cruciate ligament reconstruction. A randomized study with two-year follow-up. *Am J Sports Med* 2001;29:722–8.
- 4 Lautamies R, Härläinen A, Kettunen J, et al. Isokinetic quadriceps and hamstring muscle strength and knee function 5 years after anterior cruciate ligament reconstruction: comparison between bone-patellar tendon-bone and hamstring tendon autografts. *Knee Surg Sports Traumatol Arthrosc* 2008;16:1009–16.
- 5 Xergia SA, McClelland JA, Kvist J, et al. The influence of graft choice on isokinetic muscle strength 4–24 months after anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2011;19:768–80.
- 6 Bizzini M, Gorelick M, Munzinger U, et al. Joint laxity and isokinetic thigh muscle strength characteristics after anterior cruciate ligament reconstruction: bone patellar tendon bone versus quadrupled hamstring autografts. *Clin J Sport Med* 2006;16:4–9.
- 7 Gobbi A, Mahajan S, Zanazzo M, et al. Patellar tendon versus quadrupled bone-semi-tendinosus anterior cruciate ligament reconstruction: a prospective clinical investigation in athletes. *Arthroscopy* 2003;19:592–601.
- 8 Mattacola CG, Perrin DH, Gansneder BM, et al. Strength, functional outcome, and postural stability after anterior cruciate ligament reconstruction. *J Athl Train* 2002;37:262–8.
- 9 Jansson KA, Linko E, Sandelin J, et al. A prospective randomized study of patellar versus hamstring tendon autografts for anterior cruciate ligament reconstruction. *Am J Sports Med* 2003;31:12–18.
- 10 Bowerman SJ, Smith DR, Carlson M, et al. A comparison of factors influencing ACL injury in male and female athletes and non-athletes. *Phys Ther Sport* 2006;7:144–52.
- 11 Myer GD, Ford KR, Hewett TE. Rationale and clinical techniques for anterior cruciate ligament injury prevention among Female athletes. *J Athl Train* 2004;39:352–64.
- 12 Paterno MV, Schmitt LC, Ford KR, et al. Biomechanical measures during landing and postural stability predict second anterior cruciate ligament injury after anterior cruciate ligament reconstruction and return to sport. *Am J Sports Med* 2010;38:1968–78.
- 13 Pinczewski LA, Lyman J, Salmon LJ, et al. A 10-year comparison of anterior cruciate ligament reconstructions with hamstring tendon and patellar tendon autograft: a controlled, prospective trial. *Am J Sports Med* 2007;35:564–74.
- 14 Shelbourne KD, Gray T, Haro M. Incidence of subsequent injury to either knee within 5 years after anterior cruciate ligament reconstruction with patellar tendon autograft. *Am J Sports Med* 2009;37:246–51.
- 15 Wright RW, Dunn WR, Amendola A, et al. Risk of tearing the intact anterior cruciate ligament in the contralateral knee and rupturing the anterior cruciate ligament graft during the first 2 years after anterior cruciate ligament reconstruction: a prospective MOON cohort study. *Am J Sports Med* 2007;35:1131–4.
- 16 Barber-Westin SD, Noyes FR. Factors used to determine return to unrestricted sports activities after anterior cruciate ligament reconstruction. *Arthroscopy* 2011;27:1697–705.
- 17 Hewett TE, Di Stasi SL, Myer GD. Current concepts for injury prevention in athletes after anterior cruciate ligament reconstruction. *Am J Sports Med* 2013;41:216–24.
- 18 Lynch AD, Logerstedt DS, Grindem H, et al. Consensus criteria for defining 'successful outcome' after ACL injury and reconstruction: a Delaware-Oslo ACL cohort investigation. *Br J Sports Med* 2015;49:335–42.
- 19 van Grinsven S, van Cingel RE, Holla CJ, et al. Evidence-based rehabilitation following anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2010;18:1128–44.
- 20 de Carvalho Froufe Andrade AC, Caserotti P, de Carvalho CM, et al. Reliability of concentric, eccentric and isometric knee extension and flexion when using the REV9000 isokinetic Dynamometer. *J Human Kinet* 2013;37:47–53.
- 21 Fernandes TL, Protta TR, Fregni F, et al. Isokinetic muscle strength and knee function associated with double femoral pin fixation and fixation with interference screw in anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2012;20:275–80.
- 22 Drouin JM, Valovich-McLeod TC, Shultz SJ, et al. Reliability and validity of the Biodex system 3 pro isokinetic dynamometer velocity, torque and position measurements. *Eur J Appl Physiol* 2004;91:22–9.
- 23 Anderson JL, Lamb SE, Barker KL, et al. Changes in muscle torque following anterior cruciate ligament reconstruction: a comparison between hamstrings and patella tendon graft procedures on 45 patients. *Acta Orthop Scan* 2002;73:546–52.
- 24 Feiring DC, Ellenbecker TS, Derscheid GL. Test-retest reliability of the biodex isokinetic dynamometer. *J Orthop Sports Phys Ther* 1990;11:298–300.
- 25 Cuzzupoli S, Racette BA, Klein SE, et al. Variables associated with return to sport following anterior cruciate ligament reconstruction: a systematic review. *Br J Sports Med* 2014;48:356–64.
- 26 Hartigan EH, Axe MJ, Snyder-Mackler L. Time line for noncopers to pass return-to-sports criteria after anterior cruciate ligament reconstruction. *J Orthop Sports Phys Ther* 2010;40:141–54.
- 27 Thomee R, Kaplan Y, Kvist J, et al. Muscle strength and hop performance criteria prior to return to sports after ACL reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2011;19:1798–805.
- 28 Rosene JM, Fogarty TD, Mahaffey BL. Isokinetic hamstrings: quadriceps ratios in intercollegiate athletes. *J Athl Train* 2001;36:378–83.
- 29 Andrade MS, Cohen M, Piçarro IC, et al. Knee performance after anterior cruciate ligament reconstruction. *Isokinet Exerc Sci* 2002;10:81.
- 30 Abrams GD, Harris JD, Gupta AK, et al. Functional performance testing after anterior cruciate ligament reconstruction: a systematic review. *Orthop J Sports Med* 2014;2:1–10.
- 31 Bennell K, Wajswelner H, Lew P, et al. Isokinetic strength testing does not predict hamstring injury in Australian rules footballers. *Br J Sports Med* 1998;32:309–14.
- 32 Hewett TE, Myer GD, Zazulak BT. Hamstrings to quadriceps peak torque ratios diverge between sexes with increasing isokinetic angular velocity. *J Sci Med Sport* 2008;11:452–9.
- 33 Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ* 2009;339:b2535.
- 34 Downs SH, Black N. The feasibility of creating a checklist for the assessment of the methodological quality both of randomised and non-randomised studies of health care interventions. *J Epidemiol Community Health* 1998;52:377–84.

- 35 Ardern CL, Webster KE. Knee flexor strength recovery following hamstring tendon harvest for anterior cruciate ligament reconstruction: a systematic review. *Orthop Rev* 2009;1:e12.
- 36 Hooper P, Jutai JW, Strong G, *et al*. Age-related macular degeneration and low-vision rehabilitation: a systematic review. *Can J Ophthalmol* 2008;43:180–7.
- 37 Thomas AC, Villwock M, Wojtys EM, *et al*. Lower extremity muscle strength after anterior cruciate ligament injury and reconstruction. *J Athl Train* 2013;48: 610–20.
- 38 Beard DJ, Anderson JL, Davies S, *et al*. Hamstrings vs. patella tendon for anterior cruciate ligament reconstruction: a randomised controlled trial. *Knee* 2001;8:45–50.
- 39 Bush-Joseph CA, Hurwitz DE, Patel RR, *et al*. Dynamic function after anterior cruciate ligament reconstruction with autologous patellar tendon. *Am J Sports Med* 2001;29:36–41.
- 40 Cardone C, Menegassi Z, Emygdio R. Isokinetic assessment of muscle strength following anterior cruciate ligament reconstruction. *Isokinet Exerc Sci* 2004;12:173–7.
- 41 Carter TR, Edinger S. Isokinetic evaluation of anterior cruciate ligament reconstruction: hamstring versus patellar tendon. *Arthroscopy* 1999;15:169–72.
- 42 Eriksson K, Hamberg P, Jansson E, *et al*. Semitendinosus muscle in anterior cruciate ligament surgery: morphology and function. *Arthroscopy* 2001;17:808–17.
- 43 Feller JA, Webster KE. A randomized comparison of patellar tendon and hamstring tendon anterior cruciate ligament reconstruction. *Am J Sports Med* 2003;31:564–73.
- 44 Gokeler A, Schmalz T, Knopf E, *et al*. The relationship between isokinetic quadriceps strength and laxity on gait analysis parameters in anterior cruciate ligament reconstructed knees. *Knee Surg Sports Traumatol Arthrosc* 2003;11:372–8.
- 45 Heijne A, Werner S. A 2-year follow-up of rehabilitation after ACL reconstruction using patellar tendon or hamstring tendon grafts: a prospective randomised outcome study. *Knee Surg Sports Traumatol Arthrosc* 2010;18:805–13.
- 46 Heijne A, Werner S. Early versus late start of open kinetic chain quadriceps exercises after ACL reconstruction with patellar tendon or hamstring grafts: a prospective randomized outcome study. *Knee Surg Sports Traumatol Arthrosc* 2007;15:402–14.
- 47 Hohmann E, Tetsworth K, Bryant A. Physiotherapy-guided versus home-based, unsupervised rehabilitation in isolated anterior cruciate injuries following surgical reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2011;19:1158–67.
- 48 Janssen RP, van der Velden MJ, Pasmans HL, *et al*. Regeneration of hamstring tendons after anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2013;21:898–905.
- 49 Karanikas K, Arampatzis A, Bruggemann GP. Motor task and muscle strength followed different adaptation patterns after anterior cruciate ligament reconstruction. *Eur J Phys Rehabil Med* 2009;45:37–45.
- 50 Keays SL, Bullock-Saxton JE, Newcombe P, *et al*. The relationship between knee strength and functional stability before and after anterior cruciate ligament reconstruction. *J Orthop Res* 2003;21:231–7.
- 51 Kobayashi A, Higuchi H, Terauchi M, *et al*. Muscle performance after anterior cruciate ligament reconstruction. *Int Orthop* 2004;28:48–51.
- 52 Mikkelsen C, Werner S, Eriksson E. Closed kinetic chain alone compared to combined open and closed kinetic chain exercises for quadriceps strengthening after anterior cruciate ligament reconstruction with respect to return to sports: a prospective matched follow-up study. *Knee Surg Sports Traumatol Arthrosc* 2000;8:337–42.
- 53 Nakamura N, Horibe S, Sasaki S, *et al*. Evaluation of active knee flexion and hamstring strength after anterior cruciate ligament reconstruction using hamstring tendons. *Arthroscopy* 2002;18:598–602.
- 54 Osteras H, Augestad LB, Tondel S. Isokinetic muscle strength after anterior cruciate ligament reconstruction. *Scand J Med Sci Sports* 1998;8:279–82.
- 55 Risberg MA, Holm I, Myklebust G, *et al*. Neuromuscular training versus strength training during first 6 months after anterior cruciate ligament reconstruction: a randomized clinical trial. *Phys Ther* 2007;87:737–50.
- 56 Silva F, Ribeiro F, Oliveira J. Effect of an accelerated ACL rehabilitation protocol on knee proprioception and muscle strength after anterior cruciate ligament reconstruction. *Arch Exerc Health Dis* 2012;3:139–44.
- 57 Urabe Y, Ochi M, Onari K. Changes in isokinetic muscle strength of the lower extremity in recreational athletes with anterior cruciate ligament reconstruction. *J Sport Rehabil* 2002;11:252–67.
- 58 Webster KE, Gonzalez-Adrio R, Feller JA. Dynamic joint loading following hamstring and patellar tendon anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2004;12:15–21.
- 59 Wilk KE, Romaniello WT, Soscia SM, *et al*. The relationship between subjective knee scores, isokinetic testing, and functional testing in the ACL-reconstructed knee. *J Orthop Sports Phys Ther* 1994;20:60–73.
- 60 Witvrouw E, Bellemans J, Verdonk R, *et al*. Patellar tendon vs. doubled semitendinosus and gracilis tendon for anterior cruciate ligament reconstruction. *Int Orthop* 2001;25:308–11.
- 61 Wojtys EM, Huston LJ. Longitudinal effects of anterior cruciate ligament injury and patellar tendon autograft reconstruction on neuromuscular performance. *Am J Sports Med* 2000;28:336–44.
- 62 Xergia SA, Pappas E, Zampeli F, *et al*. Asymmetries in functional hop tests, lower extremity kinematics, and isokinetic strength persist 6 to 9 months following anterior cruciate ligament reconstruction. *J Orthop Sports Phys Ther* 2013;43:154–62.
- 63 Yoon T, Hwang J. Comparison of eccentric and concentric isokinetic exercise testing after anterior cruciate ligament reconstruction. *Yonsei Med J* 2000;41:584–92.
- 64 Yosmaoglu HB, Baltaci G, Kaya D, *et al*. Tracking ability, motor coordination, and functional determinants after anterior cruciate ligament reconstruction. *J Sport Rehabil* 2011;20:207–18.
- 65 Yosmaoglu HB, Baltaci G, Ozer H, *et al*. Effects of additional gracilis tendon harvest on muscle torque, motor coordination, and knee laxity in ACL reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2011;19:1287–92.
- 66 Yuksel HY, Erkan S, Uzun M. Factors affecting isokinetic muscle strength before and after anterior cruciate ligament reconstruction. *Acta Orthop Belg* 2011;77:339–48.
- 67 Oni OO, Crowder E. A comparison of isokinetics and muscle strength ratios following intra-articular and extra-articular reconstructions of the anterior cruciate ligament. *Injury* 1996;27:195–7.
- 68 Kellis E, Baltzopoulos V. Isokinetic eccentric exercise. *Sports Med* 1995;19:202–22.
- 69 Sole G, Hamren J, Milosavljevic S, *et al*. Test-retest reliability of isokinetic knee extension and flexion. *Arch Phys Med Rehabil* 2007;88:626–31.
- 70 Almosnino S, Stevenson JM, Bardana DD, *et al*. Reproducibility of isokinetic knee eccentric and concentric strength indices in asymptomatic young adults. *Phys Ther* 2012;13:156–62.
- 71 Li RC, Wu Y, Maffulli N, *et al*. Eccentric and concentric isokinetic knee flexion and extension: a reliability study using the Cybex 6000 dynamometer. *Br J Sports Med* 1996;30:156–60.
- 72 Baltzopoulos V, Brodie DA. Isokinetic dynamometry. Applications and limitations. *Sports Med* 1989;8:101–16.
- 73 Kannus P. Ratio of hamstring to quadriceps femoris muscles' strength in the anterior cruciate ligament insufficient knee. Relationship to long-term recovery. *Phys Ther* 1988;68:961–5.
- 74 Baltzopoulos B, King M, Gleeson N, *et al*. The BASES expert statement on measurement of muscle strength with isokinetic dynamometry. *Sport Exerc Sci* 2012:12–13.
- 75 Grimshaw P, Fowler N, Lees A, *et al*. *BIOS instant notes in sport and exercise biomechanics*. Taylor & Francis, 2004.
- 76 Gransberg L, Knutsson E. Determination of dynamic muscle strength in man with acceleration controlled isokinetic movements. *Acta Physiol Scand* 1983;119:317–20.
- 77 Maffiuletti NA, Bizzini M, Desbrosses K, *et al*. Reliability of knee extension and flexion measurements using the Con-Trex isokinetic dynamometer. *Clin Physiol Funct Imaging* 2007;27:346–53.
- 78 Greenberger HB, Wilkowsky T, Belyea B. Comparison of quadriceps peak torque using three different isokinetic dynamometers. *Isokinet Exerc Sci* 1994;4:70–5.
- 79 Heinrichs KI, Perrin DH, Weltman A, *et al*. Effect of protocol and assessment device on isokinetic peak torque of the quadriceps muscle group. *Isokinet Exerc Sci* 1995;5:7–13.



# Isokinetic muscle strength and readiness to return to sport following anterior cruciate ligament reconstruction: is there an association? A systematic review and a protocol recommendation

Marit Baste Undheim, Ciaran Cosgrave, Enda King, Siobhán Strike, Brendan Marshall, Éanna Falvey and Andrew Franklyn-Miller

*Br J Sports Med* published online June 23, 2015

---

Updated information and services can be found at:

<http://bjsm.bmj.com/content/early/2015/06/23/bjsports-2014-093962>

---

*These include:*

## Supplementary Material

Supplementary material can be found at:

<http://bjsm.bmj.com/content/suppl/2015/06/23/bjsports-2014-093962.DC1.html>

## References

This article cites 77 articles, 19 of which you can access for free at:

<http://bjsm.bmj.com/content/early/2015/06/23/bjsports-2014-093962#BIBL>

## Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

---

## Topic Collections

Articles on similar topics can be found in the following collections

[BJSM Reviews with MCQs](#) (154)  
[Procedures](#) (46)

---

## Notes

---

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>